

Concussion

How to recognise a concussion

If any of the following signs are suspected, the player should be immediately removed from play:

- Dazed look
- Motionless
- Unsteady
- Unresponsive
- Confused
- Seizure
- Clutching head
- Heightened irritability or emotion

IF IN DOUBT, SIT THEM OUT

Immediate management

Anyone with a suspected concussion should be immediately removed from play in a safe manner.

They must not return to activity that day.

If a neck injury is suspected, they should only be removed by a healthcare professional.

In all cases of suspected concussion we recommend you seek medical advice early via your GP or NHS24 111.

Return to sport, once symptom free, should be graduated. Please refer to Scottish Concussion Guidance for more information.

If any of the below suspected please ensure they are urgently taken to hospital :

- Severe neck pain
- Deteriorating consciousness
- Increasing confusion or irritability
- Severe or increasing headache
- Repeated vomiting
- Unusual behaviour change
- Seizure

- Double vision
- Weakness or tingling / burning in arms or legs

Symptoms of concussion

- Headache
- Dizziness
- Confusion
- Visual problems
- Nausea or vomiting
- Fatigue
- Drowsiness
- Pressure in head
- Sensitivity to light or noise

Questions to ask

Where are we now?

What time of day is it?

How did you get here today?

Where were you on this day last week?

For more information please visit:

<http://www.sportscotland.org.uk/concussion>

August 2019